

Incidence of Venous Thromboembolism and Benefits and Risks of Thromboprophylaxis After Cardiac Surgery: A Systematic Review and Meta-Analysis

Kwok M. Ho, MPH, PhD, FRCP; Ebrahim Bham, MBBS; Warren Pavey, MBBS

Background—Optimal thromboprophylaxis after cardiac surgery is uncertain. This systematic review aimed to define the incidence and risk factors for deep vein thrombosis (DVT), fatal and nonfatal pulmonary embolism (PE), and assess whether venous thromboembolism (VTE) prophylaxis was effective in reducing VTE without complications after cardiac surgery.

Methods and Results—Two reviewers independently searched and assessed the quality and outcomes of randomized, controlled trials (RCTs) and observational studies on VTE after cardiac surgery in the MEDLINE, EMBASE, and Cochrane controlled trial register (1966 to December 2014). Sixty-eight studies provided data on VTE outcomes or complications related to thromboprophylaxis after cardiac surgery. The majority of the studies were observational studies ($n=49$), 16 studies were RCTs, and 3 were meta-analyses. VTE prophylaxis was associated with a reduced risk of PE (relative risk [RR], 0.45; 95% confidence interval [CI], 0.28–0.72; $P=0.0008$) or symptomatic VTE (RR, 0.44; 95% CI, 0.28–0.71; $P=0.0006$) compared to the control without significant heterogeneity. Median incidence (interquartile range) of symptomatic DVT, PE, and fatal PE were 3.2% (0.6–8.1), 0.6% (0.3–2.9), and 0.3% (0.08–1.7), respectively. Previous history of VTE, obesity, left or right ventricular failure, and prolonged bed rest, mechanical ventilation, or use of a central venous catheter were common risk factors for VTE. Bleeding or cardiac tamponade requiring reoperation owing to pharmacological VTE prophylaxis alone, without systemic anticoagulation, was not observed.

Conclusions—Unless proven otherwise by adequately powered RCTs, initiating pharmacological VTE prophylaxis as soon as possible after cardiac surgery for patients who have no active bleeding is highly recommended. (*J Am Heart Assoc.* 2015;4:e002652 doi: 10.1161/JAHA.115.002652)

Key Words: deep vein thrombosis • heart surgery • prevention • pulmonary embolism

Hospitalization is a major risk factor for venous thromboembolism (VTE). According to the latest Centers for Disease Control and Prevention (CDC) analysis, approximately 547 596 hospitalizations were complicated by VTE each year among those age ≥ 18 years in the United States.¹ Deep vein thrombosis (DVT) occurred in 348 558 hospitalizations, pulmonary embolism (PE) occurred in 277 549 hospitaliza-

tions, and concomitant DVT and PE occurred in 78 511 hospitalizations each year. The total cost of VTE per person including loss of productivity and well-being was estimated to be over A\$1 470 000 in the year 2008, and it was estimated that the total cost of VTE for Australia in 2008 was A3.9 billion, more than the costs of all other diseases including cancers.²

The practice of VTE prophylaxis after cardiac surgery is controversial. According to the European Association for Cardiothoracic Surgery (EACTS) guidelines, prophylactic anticoagulation for VTE should be commenced from the first postoperative day.³ However, an increased risk of bleeding and cardiac tamponade resulting from pharmacological thromboprophylaxis remains a major concern for patients who have undergone cardiac surgery.⁴ Indeed, the latest American College of Chest Physicians evidence-based clinical practice guidelines on antithrombotic therapy and prevention of thrombosis have also stressed that patients after cardiac surgery are at high risk of major bleeding complications with only moderate risk for VTE (Grade-2C), and suggested that

From the Department of Intensive Care, Royal Perth Hospital, Perth, WA, Australia (K.M.H.); School of Population Health, University of Western Australia, Perth, WA, Australia (K.M.H.); School of Veterinary and Life Sciences, Murdoch University, Perth, WA, Australia (K.M.H., W.P.); Department of Anesthesia, Fiona Stanley Hospital, Perth, WA, Australia (E.B., W.P.).

Correspondence to: Kwok M. Ho, MPH, PhD, FRCP, FCIM, FANZCA, ICU, 4th Floor, North Block, Royal Perth Hospital, Wellington St, Perth, WA 6000, Australia. E-mail: kwok.ho@health.wa.gov.au

Received September 8, 2015; accepted September 15, 2015.

© 2015 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley Blackwell. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

pharmacological VTE prophylaxis is only needed for those with prolonged hospitalization postsurgery.⁵ These Grade-2C recommendations were mainly based on consensus among experts or weak evidence.

Recent evidence suggested that a prothrombotic state owing to multiple mechanisms, including increased fibrinogen concentrations, thrombin generation, tissue factor activation, reduced fibrinolysis, return of normal platelet aggregation, and aspirin resistance,^{6–13} is common after cardiac surgery from postoperative day 1, reaches a peak prothrombotic state between day 3 and day 5 after surgery,^{6–9} and can last up to 30 days postsurgery.⁶ Therefore, patients after cardiac surgery are not necessarily protected from VTE from day 1 postoperatively, despite their tendency for bleeding during the intraoperative and immediate postoperative period. Whether the benefits of VTE prophylaxis would outweigh its possible harms in patients after cardiac surgery remains contentious. In this systematic review, we aimed to define the incidence of DVT, fatal and nonfatal PE, risk factors for VTE, and also to assess whether VTE prophylaxis was indeed effective in reducing VTE without increasing risk of complications after cardiac surgery.

Methods

Data Sources and Study Selection

Two reviewers searched the Cochrane controlled trial register (2014, issue 4) and the EMBASE (January 1988 to December 12, 2014) and MEDLINE databases (1966 to December 12, 2014) independently. During the electronic database search, the following exploded Medical Subject Heading (MeSH) terms were used: “heart surgery” or “cardiac surgery” or “coronary artery bypass” or “valve surgery” or “valvular surgery” AND “venous thromboembolism” or “pulmonary embolism” or “deep vein thrombosis”. The search included all forms of publications, including case series, cohort or case control studies, clinical trials, letters, editorials, reviews, meta-analysis, or randomized, controlled trials (RCTs) without any age or language restrictions. Animal studies, studies without involving cardiac surgery, or isolated case reports were excluded from this review. The reference lists of related editorials, reviews, and original articles identified were searched for relevant studies. Finally, the websites of the International Network of Agencies of Health Technology Assessment in Health Care were searched to ensure all suitable trials were included.

Two reviewers independently examined all identified studies to confirm they fulfilled the inclusion criteria and VTE outcomes were reported. For RCTs, the quality of the trial, such as allocation concealment, randomization method, and blinding of treatment and assessment of outcome, was assessed. The grading of allocation was based on the

Cochrane approach, that is, adequate, inadequate, or uncertain. When the reported methodology and results of the included trials were unclear, the corresponding authors of the trials were contacted to clarify the data ($n=2$, but 1 did not respond to our study inquiries). Any disagreements between the 2 independent reviewers were resolved by consensus.

Data Reporting and Statistical Analysis

VTE was the primary outcome of this review, and, whenever possible, we reported the incidence of symptomatic and asymptomatic VTE separately. In trying to identify the risk factors for VTE and complications after the use of VTE prophylaxis, only risk factors that were significant in the multivariate analyses were reported in this review. We also assessed whether the incidence of DVT was different between the leg with and without the great saphenous vein harvested for coronary artery bypass grafting (CABG) surgery.

The other outcomes assessed included the risk factors for bleeding complications or cardiac tamponade. In trying to identify whether VTE prophylaxis, in general, would be effective in reducing VTE after cardiac surgery, all interventions aimed to VTE were analyzed together against no comparative VTE prophylaxis. Outcomes were reported as relative risk (RR) with 95% confidence interval (CI), using a random-effects model (with the Mantel-Haenszel method, which estimates the amount of between-study variation by comparing each study's result with a Mantel-Haenszel fixed-effect meta-analysis result). The presence of heterogeneity between trials was assessed by the chi-square statistics and the extent of inconsistency was assessed using I^2 statistics.¹⁴ An $I^2 > 40\%$ was considered as significant heterogeneity in this study, and when heterogeneity was present, the data were further examined to explore for reasons for the heterogeneity by metaregression. Data were analyzed by the Review Manager (version 4.2.6 for Windows; Oxford, UK: The Cochrane Collaboration, 2003) and metaregression was conducted by Comprehensive Meta-analysis (version 2.2.034, 2006, USA). A P value of less than 0.05 was taken as significant in this systematic review.

Sensitivity Analysis

In assessing the incidence of VTE, we did a sensitivity analysis by restricting the analysis to higher-quality studies (RCTs and prospective cohort studies). We also assessed whether the incidence of VTE was related to the sample size of the studies and how recent the study was conducted. In assessing effectiveness of VTE prophylaxis, we did a sensitivity analysis by restricting our analysis to studies that had adequate allocation concealment. In assessing whether the leg with the great saphenous vein harvested was associated with a higher




Figure 1. Flowchart showing the inclusion and exclusion of studies for this systematic review.

risk of DVT, we restricted our analysis to studies that utilized routine ultrasound scans on both legs regardless of whether the patients had symptoms of DVT.

Results

Study Characteristics

Of the 6636 studies identified in the literature search, a total of 68 studies provided data on VTE outcomes or complications related to VTE prophylaxis after cardiac surgery (Figure 1). The majority of the studies were observational studies ($n=49$), 16 studies were RCTs, and 3 were meta-analyses (Tables 1 and 2).^{15–84} Of the 16 RCTs identified,* only 9 studies had allocation concealment and 6 used some form of blinding (Table 1). Seven RCTs assessed 1 method of VTE prophylaxis against a placebo or control group, and 5 studies compared 1 pharmacological agent against another

(eg, 3 studies on warfarin vs. aspirin, 2 studies on unfractionated heparin [UFH] vs. low-molecular-weight heparin [LMWH] or hirudin). Six RCTs compared interventions primarily not aiming at reducing VTE, but VTE data were reported (eg, activated factor VIIa, off-pump CABG vs. on-pump CABG, placebo vs. cyclooxygenase 2 inhibitors, and different types of autograft/prosthesis or cell saver devices) (Table 1). The 3 meta-analyses identified were related to use of factor VIIa or tranexamic acid for adult cardiac surgery and warfarin versus antiplatelet agents after Fontan surgery for congenital heart diseases.^{62,63,74}

Of the 49 observational studies included in this study, only 9 (18%) were prospective studies and 13 reported risk factors for VTE events or complications from VTE prophylaxis after adjusting for known risk factors for VTE or complications with a multivariate analysis, respectively (Table 2).

VTE Events After Cardiac Surgery With Prophylaxis Versus Control/Placebo

Pooling all forms of VTE prophylaxis after cardiac surgery together, VTE prophylaxis was associated with a reduced risk of PE (RR, 0.45; 95% CI, 0.28–0.72; $P=0.0008$; $I^2=0\%$) and symptomatic VTE (RR, 0.44; 95% CI, 0.28–0.71; $P=0.0006$; $I^2=0\%$) (Figures 2 and 3) compared to the control group without significant heterogeneity. After restricting our analyses to only studies that had adequate allocation concealment,** the benefits of VTE prophylaxis on risks of PE (RR, 0.38; 95% CI, 0.23–0.63; $I^2=0\%$) and symptomatic VTE (RR, 0.38; 95% CI, 0.23–0.62; $I^2=0\%$) remained unchanged (Figures 4 and 5, respectively). Similarly, restricting the analysis to adult cardiac surgical patients only, the benefits of VTE prophylaxis on risks of PE (RR, 0.45; 95% CI, 0.28–0.72; $I^2=0\%$) and symptomatic VTE also remained unchanged (RR, 0.44; 95% CI, 0.28–0.67; $I^2=0\%$). Using VTE as an outcome measure, the funnel plot did not suggest presence of a significant publication bias (Figure 6).

VTE Events After 2 Different VTE Pharmacological Prophylaxis Agents

Only a small number of patients ($n=276$) had been enrolled in 3 RCTs comparing warfarin against high-dose aspirin (>300 mg/day for adults, 5 mg/kg per day for children), and the risk of VTE was not different (RR, 0.97; 95% CI, 0.50–1.89; $P=0.94$; $I^2=0\%$). One meta-analysis comparing aspirin with warfarin after Fontan surgery also did not show a significant difference in symptomatic VTE.⁶³ Similarly, only a very small number of patients ($n=59$) had been enrolled in a direct comparison between UFH and LMWH or hirudin, and the risk of VTE was also not different between the 2 groups

*References 16, 17, 33, 36, 39, 42, 45, 52, 55, 56, 64, 66, 68–70, 78.

**References 16, 17, 36, 39, 42, 45, 55, 68, 69.

Table 1. Characteristics of the Included RCT and Meta-Analysis Comparing Different Interventions With Outcome Data on VTE

Study (Year of Publication, First Author, Reference Number)	Type of Surgery and Sample Size	Mean Age, y (or No. of Studies in Meta-Analysis)	Interventions	Outcomes Including Complications	Bias Assessment
1996, Ramos et al. [16]	Mixed (n=2551)	64	IPC stockings till fully ambulatory, both groups received UFH	Symptomatic PE (4% control vs. 1.5% IPC), complications not reported	Allocation concealment adequate, no blinding, loss to follow-up (0%)
2013, Mirhosseini et al. [17]	Off-pump CABG (n=120)	62	UFH alone vs UFH+aspirin (80 mg daily) for the first 7 to 10 days	Asymptomatic lower-limb DVT (16.6% UFH alone vs 3.3% UFH+aspirin). Postoperative bleeding (1.7% UFH alone vs. 6.7% UFH+aspirin, no multivariate analysis was reported)	Allocation concealment adequate, patients and research nurses blinded to the intervention, loss to follow-up (0%)
2013, Ayatollahzadeh-Isfahani et al. [36]	CABG (n=185)	60	Leg elevation during surgery vs. supine position, no UFH was used for both groups	Asymptomatic lower-limb DVT (18.4% supine group vs. 8.6% leg elevation group; ipsilateral leg 21/185 vs. contralateral leg 4/185)	Allocation concealment adequate, no blinding, loss to follow-up (0%)
2005, Nussmeier et al. [39]	On-pump CABG (n=1636)	62	Placebo vs. valdecoxib or valdecoxib+parecoxib. All patients received aspirin and standard VTE prophylaxis for all 3 groups	Symptomatic PE (0.2% placebo vs. 0.4% for both valdecoxib and valdecoxib+parecoxib groups)	Allocation concealment adequate, double-blinded, loss to follow-up (2%)
2009, Gill et al. [42]	Mixed (n=172)	64	Placebo vs. activated factor VIIa (40 µg/kg) vs. activate factor VIIa (80 µg/kg)	Symptomatic PE (0% for all 3 groups)	Allocation concealment adequate, double-blinded, loss to follow-up (3.9%)
2013, McCrindle et al. [45]	Fontan procedure (n=111)	5	Aspirin vs. warfarin	Symptomatic VTE (7% aspirin vs. 5.8% warfarin) and asymptomatic VTE (14% aspirin vs. 19% warfarin)	Allocation concealment adequate, single-blinded, loss to follow-up (2.7%)
2011, Doss et al. [52]	Aortic valve replacement (n=60)	62	Pulmonary autograft vs. mechanical valve, stentless bioprosthesis vs. mechanical valve, or stentless vs. stented bioprosthesis	Symptomatic PE (5% in the stented bioprosthetic group, 0% for the all those 5 groups)	Allocation concealment uncertain, no blinding, loss to follow-up (0%)
2003, Lee et al. [55]	CABG (n=60)	66	Off-pump vs on-pump CABG	Symptomatic PE (3.3% in the off-pump group vs. 0% in the on-pump group)	Allocation concealment adequate, single-blinded, loss to follow-up (0%)
2007, Riess et al. [56]	CABG (n=20)	58	IV lepirudin vs. IV heparin on bypass and first 2 days after surgery	Symptomatic PE (10% in the lepirudin group vs. 0% for the IV heparin group)	Allocation concealment unclear, no blinding, loss to follow-up (0%)
2011, Adler Ma et al. [62]	Off-pump CABG (n=544)	8 studies	IV tranexamic acid for patients undergoing off-pump CABG	Symptomatic VTE (0.7% tranexamic acid vs. 1.5% control group)	Allocation concealment unclear in 25% and also no blinding in 25% of the studies. No significant heterogeneity on VTE data between studies
2011, Marrone et al. [63]	Fontan surgery (n=1075)	20 studies	Warfarin vs. antiplatelet agents	Symptomatic VTE (5% anticoagulation vs. 4.5% antiplatelet agents)	95% were observational studies. No significant heterogeneity on VTE data between studies

Continued

Table 1. Continued

Study (Year of Publication, First Author, Reference Number)	Type of Surgery and Sample Size	Mean Age, y (or No. of Studies in Meta-Analysis)	Interventions	Outcomes Including Complications		Bias Assessment
					Symptomatic VTE (0 vs. 0%) by USS	
1993, Beghi et al. [64]	Mixed (n=39)	60	LMWH-Fluxum 3200 anti-Xa units daily vs. calcium heparin 5000 IU 3 times daily			Allocation concealment unclear, no blinding, loss to follow-up (0%)
1995, Goldhaber et al. [66]	CABG (n=344)	64	IPC with graduated compression stockings vs. graduated compression stockings alone. All patients received aspirin		Asymptomatic DVT before discharge (19% in IPC vs. 22% in control group). Symptomatic PE (1.2% in IPC vs. 0.6% in control group)	Allocation concealment unclear, no blinding, loss to follow-up (4%)
1979, Pantely et al. [68]	CABG (n=65)	54	Warfarin vs control vs. high-dose antiplatelet agent group (aspirin 325 mg+dipyridamole 75 mg both 3 times per day)		Symptomatic PE (6.7% in control, 0% in antiplatelet agent group and warfarin group). Symptomatic DVT (3.3% in control, 0% in antiplatelet agent and warfarin group)	Allocation concealment adequate, no blinding, loss to follow-up to VTE outcomes (0%)
2010, Schroeder et al. [69]	Congenital heart diseases (n=90)	4 months	IV heparin (10 U/kg per hour vs. 5% dextrose) for 14 days or until central venous catheter removed		Asymptomatic central venous catheter related thrombosis (15% heparin vs. 16% placebo group)	Allocation concealment adequate, double-blinded, loss to follow-up (0%)
2013, Weitert et al. [70]	Mixed (n=1049)	67	CardioPAT intra- and postoperative cell saver system vs. standard intraoperative cell saver system (hemonetics) without using postoperative cell saver		Symptomatic DVT (1.9% CardioPAT vs. 2.7% hemonetics system)	Allocation concealment unclear, no blinding, loss to follow-up (0%)
2011, Ponschab et al. [74]	Mixed (n=470)	6 studies including the study by Gill [28]	Recombinant activated factor VII vs. placebo		Outcomes on symptomatic VTE extracted from the studies included in this meta-analysis (0.4% factor VIIa vs. 0% control)	Only 2 studies were double-blinded RCTs with adequate allocation concealment, no significant heterogeneity between studies on VTE outcomes
1982, McEnany et al. [78]	CABG (n=216)	50	Placebo vs. aspirin (300 mg bd) vs. warfarin (prothrombin time 1.5 to 2.0 of control)		Symptomatic PE (1.3% placebo vs. 2.8% aspirin vs. 1.5% warfarin)	Allocation concealment unclear, partial blinding on warfarin group and double-blinded for aspirin group, loss to follow-up (0%)
2006, Lahtiainen et al. [33]	Off-pump CABG, no UFH unless in atrial fibrillation (12.5% of the included patients) (n=24)	67	Secondary analysis of an RCT comparing a new proximal aortic anastomotic device to a conventional hand-sewn technique		Routine CTPA 1 week after surgery (all asymptomatic, PE 25% in each treatment arm)	Allocation concealment unclear, unblinded, loss to follow-up (0%) (but the study was terminated prematurely owing to unavailability of the anastomotic devices)

Mixed surgery includes CABG, heart valve, and other types of cardiac surgery. CABG indicates coronary artery bypass grafting; CTPA, computed tomographic pulmonary angiogram; DVT, deep vein thrombosis; IPC, intermittent pneumatic compression; LMWH, low-molecular-weight heparin; PE, pulmonary embolism; RCT, randomized, controlled trials; UFH, unfractionated heparin; USS, ultrasound scan; VTE, venous thromboembolism.

Table 2. Characteristics of Original Studies Reporting on Incidence and Risk Factors for VTE After Cardiac Surgery

Study (Year of Publication, First Author, [Reference Number])	Type of Study and Use of LMWH, UFH, or Other Forms of VTE Prophylaxis	Type of Surgery and Sample Size	Duration of Follow-up	Incidence of Symptomatic DVT and PE, Diagnostic Modality	Incidence of Asymptomatic DVT and PE (%), Diagnostic Modality	Significant Adjusted Risk Factors for VTE and Complications	Bias Assessment
1995, Ramos et al. [15], likely same cohort of patients as in [16]	Cohort, UFH 5000 U bd	Mixed (n=2551)	30 days	PE (2.7%) by pulmonary angiography or high-probability V/Q scan, fatal PE (0.08%) by postmortem	Not reported	VTE: 1. Prior history of VTE (OR, 3.1). 2. Obesity (OR, 2.6). 3. Preoperative LVEF <40% (OR, 6.8). 4. Previous right heart catheterization (OR, 2.9). 5. IPPV >3 days (OR, 2.5). 6. HITSS (OR, 2.5). 7. Postoperative CHF (OR, 4.1)	Retrospective, no blinding
2013, Mirhosseini et al. [17]	RCT, UFH 5000 U tds	Off-pump CABG (n=120)	Uncertain	PE (0%), unclear modality	DVT (10%) by USS	No multivariate analysis was conducted, bleeding of uncertain severity occurred in 4 patients (6.7%) after UFH with aspirin against 1 patient (1.7%) with heparin alone ($P=0.34$).	Prospective, double-blinded
1999, Pouplard et al. [18]	Cohort, UFH 120 U/kg per day followed by 200 U/kg per day after valve surgery, LMWH after CABG	Valve (n=157)+CABG (n=150)	10 days	Fatal PE (0.3%) (in a patient with HIT)	Not reported	No multivariate analysis was conducted.	Prospective, no blinding
2012, Kulik et al. [19]	Cohort, 5 different VTE prophylaxis (1) mechanical only (2) LMWH or UFH (3) fondaparinux (4) mechanical+LMWH or UFH (5) no prophylaxis, all these strategies were measured at 48 hours after surgery	Either off (19%) or on-pump (91%) CABG (n=92 699)	6 weeks	VTE (0.74%) by USS or venography, or CTPA or V/Q scan	Not reported	VTE: No difference between the 5 strategies of VTE prophylaxis (HR, 1.07 to 1.16 compared to no VTE prophylaxis) Bleeding: No difference between the 5 strategies of VTE prophylaxis (HR, 0.9 to 1.01 compared to no VTE prophylaxis: 1.5%, 1.7%, 3.4%, 1.5%, 1.4%, respectively)	Retrospective, no blinding

Continued

Table 2. Continued

Study (Year of Publication, First Author, [Reference Number])	Type of Study and Use of LMWH, UFH, or Other Forms of VTE Prophylaxis	Type of Surgery and Sample Size	Duration of Follow-up	Incidence of Symptomatic DVT and PE, Diagnostic Modality	Incidence of Asymptomatic DVT and PE (%), Diagnostic Modality	Significant Adjusted Risk Factors for VTE and Complications	Bias Assessment
2004, Ambrosetti et al. [20,27]	Cohort, UFH not used in 37% or used <3 days in 38%, bilateral GCS used only in 7%	CABG (n=270)	4 to 19 days after surgery (on admission to rehabilitation center)	Fatal PE (0.4%) and PE (0.7%) by V/Q scan	DVT (17.4%): 23/47 on the leg without the great saphenous vein harvested vs. 24/47 on the leg with the vein harvested; 21% for those without UFH by routine USS screening	VTE: Age, female, postoperative complications, and no UFH before discharge from the surgery unit were predictive of DVT on admission to rehabilitation unit (all $P<0.001$, but ORs not reported). Cardiac tamponade: Routine echocardiography (n=235) at time of USS and then 2 and 4 weeks later: no differences in incidence of tamponade between those with and without systemic anticoagulation	Prospective, no blinding
2008, Frizzelli et al. [21]	Cohort, UFH 200 U/kg per day after removal of drains for 48 to 72 hours after CABG, LMWH followed by warfarin after prosthetic valve surgery or in atrial fibrillation after CABG	Mixed (n=815)	5 to 7 days after surgery (on admission to rehabilitation center)	PE (0.7%) by CTPA: 3.5% for those on aspirin+UFH vs. 0% for those on warfarin+UFH	Central venous catheter related DVT (48%) routine USS	No multivariate analysis was conducted.	Prospective, no blinding
1991, Delaria and Hunter [22]	Cohort, no routine UFH was used after CABG, warfarin after valve surgery	Mixed (75% CABG) (n=10 638)	Uncertain	DVT (0.7%) (20/36 on the leg without the great saphenous vein harvested vs. 18/36 on the leg with great saphenous vein harvested) by USS or venography, PE (0.4%) by high probability V/Q scan or pulmonary angiography, fatal PE (0.09%)	Not reported	No multivariate analysis was conducted.	Retrospective, no blinding
2013, Ho et al. [23]	Cohort, routine UFH for all patients	Mixed (n=2131)	30 months	VTE (1.6%), fatal PE (0.2%), diagnosis by ICD-10 codes	Not reported	VTE: 1. Body mass index (OR, 1.2 per index increment). 2. Charlson comorbidity index (OR, 1.20 per index increment)	Retrospective, no blinding

Continued

Table 2. Continued

Study (Year of Publication, First Author, [Reference Number])	Type of Study and Use of LMWH, UFH, or Other Forms of VTE Prophylaxis	Type of Surgery and Sample Size	Duration of Follow-up	Incidence of Symptomatic DVT and PE, Diagnostic Modality	Incidence of Asymptomatic DVT and PE (%), Diagnostic Modality	Significant Adjusted Risk Factors for VTE and Complications	Bias Assessment
1982, Hanson and Levine [24]	Cohort, no routine UFH and antiplatelet agents	CABG (n=5000)	Uncertain	PE (0.5%) by high probability V/Q scan or pulmonary angiography, fatal PE (0.06%)	Not reported	No multivariate analysis was conducted.	Retrospective, no blinding
2010, Schwann et al. [25]	Cohort, routine dual antiplatelet agents after CABG, aspirin+warfarin for combined valve-CABG, warfarin for valve alone, all patients also received enoxaparin 40 mg/day (if <100 kg)+bilateral intermittent pneumatic compression+daily ambulation	Mixed (n=1070)	Day 3 to 4 first USS and then weekly until hospital discharge	PE (0.09%)	DVT (13%) (62/139 on the leg without the great saphenous vein harvested vs. 114/139 on the leg with the great saphenous vein harvested) by routine USS on day 3 to 4 and then day 6 to 7 and then weekly	VTE: 1. Age (OR, 1.2 per 10 year increment). 2. Reintubation (OR, 2.57). 3. Transfusion (OR, 2.24). 4. Ventilation (OR, 1.02 per 10 hours) Bleeding requiring re-exploration was not different between those with DVT treated with systemic anticoagulation and those without DVT.	Prospective, no blinding
2013, Saranteas et al. [26]	Cohort, pharmacological and mechanical VTE prophylaxis	Mixed (n=315)	24 hours after surgery	DVT (0.6%) by unclear modality, fatal PE (0.3%)	Inferior vena cava thrombus (2.5%) by echocardiography	No multivariate analysis was conducted.	Retrospective, no blinding
1975, Rao et al. [28]	Cohort, GCS and ambulation only, no UFH	CABG (n=231)	Uncertain	PE (9.5%) by high probability V/Q scan or pulmonary angiography, fatal PE (1.7%)	Not reported	No multivariate analysis was conducted.	Retrospective, no blinding
1993, Josa et al. [29]	Cohort, warfarin from day 3 after heart valve surgery, no UFH postoperatively	Mixed (valve: n=120; CABG: 819; CABG+valve: n=94) (total 1033)	Uncertain (but PE was reported up to 2 weeks)	PE (overall 3.2%, 3.9% after CABG, 0% after valve, 1% after combined CABG+valve) by high probability V/Q scan or pulmonary angiography, fatal PE (0.6%) by postmortem, DVT (0.7%) (1/819 on the contralateral to harvest side vs. 5/819 on the harvest side) was investigated only after PE was diagnosed	Not reported	No multivariate analysis was conducted.	Retrospective, no blinding

Continued

Table 2. Continued

Study (Year of Publication, First Author, [Reference Number])	Type of Study and Use of LMWH, UFH, or Other Forms of VTE Prophylaxis	Type of Surgery and Sample Size	Duration of Follow-up	Incidence of Symptomatic DVT and PE, Diagnostic Modality	Incidence of Asymptomatic DVT and PE, Diagnostic Modality	Significant Adjusted Risk Factors for VTE and Complications	Bias Assessment
1992, Gillinov et al. [30]	Case-control, warfarin after valve surgery, aspirin and dipyridamole after CABG	Mixed (n=64)	Uncertain	PE (0.6%) by high probability V/Q scan or CTPA, fatal PE (0.2%) by postmortem	Not reported	VTE: 1. Preoperative bed rest (OR, 6). 2. >Hospital day preoperatively (OR, 6). 3. <15 day between coronary catheterization and surgery (OR, 5). 4. Postoperative congestive heart failure (OR, 8). 5. Postoperative bed rest >3 days (OR, 5)	Retrospective, no blinding
1991, Reis et al. [31]	Cohort (nonconsecutive), GCS and aspirin only, no UFH	CABG (n=29)	Until hospital discharge	Not reported	DVT (48%): (10/29 on the leg without the great saphenous vein harvested vs. 10/29 on the leg with the vein harvested), routinely on both legs for all patients	No multivariate analysis was conducted.	Prospective, no blinding
2009, Egawa et al. [32]	Two cohort: (1) No VTE guidelines, (2) GCS for all, intermittent pneumatic compression for moderate to high-risk patients, anticoagulation for high-risk patients	Mixed (1st group: 1467, 2nd group: 1389)	Until hospital discharge (mean 22 to 25 days)	Group 1: PE (0.4%), Group 2: PE (0%), CTPA	Not reported	No multivariate analysis was conducted.	Retrospective, no blinding
2006, Lahtinen et al. [33]	Secondary analysis of a RCT comparing a new proximal aortic anastomotic device to a conventional hand-sewn technique, no UFH unless in atrial fibrillation (12.5% of the included patients)	Off-pump CABG (n=24)	1 week after surgery	PE (0%)	PE (25%) by routine CTPA	No multivariate analysis was conducted.	Prospective, no blinding

Continued

Table 2. Continued

Study (Year of Publication, First Author, [Reference Number])	Type of Study and Use of LMWH, UFH, or Other Forms of VTE Prophylaxis	Type of Surgery and Sample Size	Duration of Follow-up	Incidence of Symptomatic DVT and PE, Diagnostic Modality	Incidence of Asymptomatic DVT and PE (%), Diagnostic Modality	Significant Adjusted Risk Factors for VTE and Complications	Bias Assessment
2010, Mitra et al. [34]	Two cohorts (1st group: received activated factor VIIa, 2nd group: cardiothoracic registry), VTE prophylaxis strategies uncertain	Mixed (1st group: n=705, 2nd group: n=6554)	Uncertain	PE (1st group: 0.4%, 2nd group: 7%), unclear modality	Not reported	VTE: 1. Urgent surgery (OR, 2.8). 2. Massive transfusion (OR, 2.8). 3. Activated factor VIIa (OR, 0.02)	Retrospective, no blinding
1992, Canver and Fiedler [35]	Cohort, no UFH	Mixed (n=4393)	Until hospital discharge	PE (0.1%), fatal PE (0.02%), DVT (0.2%), venous Doppler, contrast venography, lung isotope scan or pulmonary angiography	Not reported	No multivariate analysis was conducted, 1 patient developed HIT and retroperitoneal hematoma after initiating systemic UFH with DVT	Retrospective, no blinding
2013, Ayatollahzadeh-Isfahani et al. [36]	RCT, no UFH	CABG (n=185)	Until 5 to 7 days after surgery	DVT (0.5%) by USS	DVT (13.5%) by routine USS	No multivariate analysis result was reported.	Prospective, no blinding
2007, Gandhi et al. [37]	Cohort, VTE prophylaxis strategies uncertain	Ventricular assist device or orthotopic heart transplant requiring activated factor VIIa (n=15)	Until hospital discharge	DVT (6.7%), unclear modality	Not reported	No multivariate analysis was conducted.	Retrospective, no blinding
2007, Mueller et al. [38]	Cohort, VTE prophylaxis strategies uncertain	CABG (n=259)	5 days after surgery	Not reported	PE (1.9%) by CTPA	No multivariate analysis was conducted.	Prospective, no blinding
2005, Nussmeier et al. [39]	RCT, aspirin for all patients and standard VTE prophylaxis strategy permitted	On-pump CABG (n=1636)	30 days after surgery	PE (0.3%), DVT (0.06%), unclear modality	Not reported	No multivariate analysis on VTE was conducted.	Prospective, double-blinded
2009, Belczak et al. [40]	Cohort (nonconsecutive), VTE prophylaxis strategies uncertain	CABG (n=44)	90 days after surgery	DVT (2.3%) on the leg with long-saphenous vein harvested, USS modality	Not reported	No multivariate analysis was conducted.	Retrospective, no blinding
2011, ElBardissi et al. [41]	Cohort, VTE prophylaxis strategies uncertain	Transcatheter aortic valve implantation (n=249)	30 days	PE (0.4%), unclear modality	Not reported	No multivariate analysis was conducted.	Retrospective, no blinding

Continued

Table 2. Continued

Study (Year of Publication, First Author, [Reference Number])	Type of Study and Use of LMWH, UFH, or Other Forms of VTE Prophylaxis	Type of Surgery and Sample Size	Duration of Follow-up	Incidence of Symptomatic DVT and PE, Diagnostic Modality	Incidence of Asymptomatic DVT and PE (%), Diagnostic Modality	Significant Adjusted Risk Factors for VTE and Complications	Bias Assessment
2009, Gill et al. [42]	RCT, VTE prophylaxis uncertain	Mixed (n=172)	30 days	PE (0%), clinical signs with postmortem or V/Q scan, DVT by venography or duplex USS	Not reported	No multivariate analysis was conducted.	Prospective, double-blinded
2014, Alfrevic et al. [43]	Case-control, VTE prophylaxis strategies uncertain	Mixed (n=303)	Until hospital discharge	VTE (9.5%), unclear modality	Not reported	No multivariate analysis was conducted.	Retrospective, no blinding
2003, White et al. [44]	Cohort, VTE prophylaxis strategies uncertain	Mixed. Group 1: without malignancy (valve: n=16, 0.36, CABG: n=66, 1.80). Group 2: with malignancy+CABG (n=2243)	90 days	Without malignancy: VTE (heart valve 0.5% [0.2% after hospital discharge], CABG 1.1% [0.7% after hospital discharge]). With malignancy: VTE (CABG 1.5% with 0.8% after discharge), VTE by ICD-9 codes, unclear modality	Not reported	No multivariate analysis was conducted.	Retrospective, no blinding
2013, McCrindle et al. [45]	RCT, aspirin (5 mg/kg per day) vs. warfarin (INR 2 to 3) as VTE prophylaxis	Fontan surgery (n=111)	2.5 years	VTE (17%), TEE or echocardiography	VTE (6.5%), routine TEE or echocardiography	VTE: 1. Uncontrolled warfarin, INR between 2 and 3 <30% time (OR, 3.5). 2. Central venous catheter >10 days or at ICU discharge (OR, 17.8). 3. Pulmonary atresia with intact ventricular septum (OR, 3.6). 4. Lower preoperative bilirubin (OR, 0.8).	Prospective, single-blinded
2013, Zahn et al. [46]	Cohort, VTE prophylaxis strategies uncertain	Transcatheter aortic valve implantation (n=1318)	12 months	PE (1.6%), unclear modality	Not reported	No multivariate analysis on VTE was conducted.	Retrospective, no blinding
2012, Iribarne et al. [47]	Cohort, VTE prophylaxis strategies uncertain	Isolated mitral valve surgery (n=6297)	Until hospital discharge	VTE (0.8%), unclear modality	Not reported	No multivariate analysis on VTE was conducted.	Retrospective, no blinding

Continued

Table 2. Continued

Study (Year of Publication, First Author, [Reference Number])	Type of Study and Use of LMWH, UFH, or Other Forms of VTE Prophylaxis	Type of Surgery and Sample Size	Duration of Follow-up	Incidence of Symptomatic DVT and PE, Diagnostic Modality	Incidence of Asymptomatic DVT and PE (%), Diagnostic Modality	Significant Adjusted Risk Factors for VTE and Complications	Bias Assessment
2010, Thielmann et al. [48]	Cohort, VTE prophylaxis strategies uncertain	Mixed (n=153), including patients who had a low platelet count and an HIT test was initiated	Until hospital discharge	PE (3.3%), unclear modality	Not reported	No multivariate analysis on VTE was conducted.	Retrospective, no blinding
1995, Rosenthal et al. [49]	Cohort, no UFH, aspirin or warfarin	Fontan surgery (n=70)	1.9 to 5.5 years	VTE (20%), TEE or CT PA	Not reported	No multivariate analysis on VTE was conducted.	Retrospective, no blinding
1999, Michalopoulos et al. [50]	Case-control, VTE prophylaxis strategies uncertain	CABG (n=2014)	Until hospital discharge	Fatal PE (0.05%), unclear modality	Not reported	No multivariate analysis on VTE was conducted.	Retrospective, no blinding
1996, Briffa and Large [51]	Cohort, VTE prophylaxis strategies uncertain	CABG (n=1400)	Until hospital discharge	PE (overall 0.5%) (urgent bed rest before surgery cases 2.2%, nonurgent cases 0.2%), unclear modality	Not reported	No multivariate analysis on VTE was conducted.	Retrospective, no blinding
2011, Doss et al. [52]	RCT, warfarin for 3 months after bioprosthetic, lifelong warfarin after mechanical valve, and no anticoagulation with autografts	Aortic valve replacement (n=120)	12 months	PE (0.8%)	Not reported	No multivariate analysis on VTE was conducted, 3 patients (2.5%) treated with warfarin had bleeding requiring reoperation.	Prospective, no blinding
2011, Bucci et al. [53]	Cohort, UFH (SC for aortic mechanical or tissue valve or mitral tissue valve; IV UFH for mechanical mitral valve) or LMWH (double dose for mechanical mitral compared to tissue mitral or all types of aortic) until INR within targets	Valve (n=203)	Until hospital discharge	PE (0%), unclear modality	Not reported	No multivariate analysis on VTE was conducted. Major bleeding (6.4%). Systemic anticoagulation with concomitant aspirin and/or clopidogrel (OR, 7.4), 6 patients (3%) had international normalized ratio >5 postoperatively.	Prospective, no blinding
2000, Montalescot et al. [54]	Cohort, systemic anticoagulation IV UFH or SC LMWH from day 6 after surgery	Single or double mechanical valve surgery (n=208)	Until hospital discharge	PE (0%), unclear modality	Not reported	No multivariate analysis on VTE was conducted. Bleeding occurred in 4 patients (1.9%) (2 after UFH and 2 after LMWH) with 1 patient with UFH being systemically overanticoagulated	Retrospective, no blinding

Continued

Table 2. Continued

Study (Year of Publication, First Author, [Reference Number])	Type of Study and Use of LMWH, UFH, or Other Forms of VTE Prophylaxis	Type of Surgery and Sample Size	Duration of Follow-up	Incidence of Symptomatic DVT and PE, Diagnostic Modality	Incidence of Asymptomatic DVT and PE (%), Diagnostic Modality	Significant Adjusted Risk Factors for VTE and Complications	Bias Assessment
2003, Lee et al. [55]	RCT, on-pump vs. off-pump, VTE prophylaxis strategies uncertain	CABG (n=60)	Until hospital discharge	Fatal PE (1.7%), unclear modality	Not reported	No multivariate analysis on VTE was conducted.	Prospective, single-blinded
2007, Riess et al. [56]	RCT, IV lepirudin vs. IV heparin on bypass and first 2 days after surgery	CABG (n=20)	Until day 3 after surgery	PE (5%), unclear modality	Not reported	No multivariate analysis on VTE was conducted.	Prospective, no blinding
2005, Rastan et al. [57]	Cohort (autopsy study), VTE prophylaxis strategies uncertain	Mixed (n=468)	Mean survival time 14 days after surgery	Fatal PE (total 6.6%, 3.8% not diagnosed before postmortem accounting for 11.4% of the unexplained deaths)	Not reported	No multivariate analysis on VTE was conducted.	Retrospective, no blinding
1997, Zehr et al. [58]	Cohort (autopsy study), VTE prophylaxis strategies uncertain	Mixed (n=147)	Mean survival time 22 days after surgery	Fatal PE (total 4.1%, 1.4% not diagnosed before postmortem accounting for 20% of the unexplained deaths)	Not reported	No multivariate analysis on VTE was conducted.	Retrospective, no blinding
1976, Heard [59]	Cohort, VTE prophylaxis strategies uncertain	CABG (n=80)	13 months	Fatal PE (2.5%), diagnosed by postmortem	Not reported	No multivariate analysis on VTE was conducted.	Retrospective, no blinding
2014, Song et al. [60]	Cohort, VTE prophylaxis strategies uncertain	Mixed (n=25) who received prothrombin complex concentrates for coagulopathy	Until hospital discharge	Central venous catheter related DVT (4%), unclear modality	Not reported	No multivariate analysis on VTE was conducted.	Retrospective, no blinding
1994, Sanchez and Haft [61]	Cohort, VTE prophylaxis strategies uncertain	CABG (n=224)	30 days	PE (0.5%) by V/Q scan, DVT (4%) by clinical criteria with a negative V/Q scan	Not reported	No multivariate analysis on VTE was conducted.	Retrospective, no blinding
2013, Clark et al. [65]	Cohort, VTE prophylaxis strategies uncertain	Mixed (n=11) and received factor IX complex for bleeding	Until hospital discharge	PE (9.1%)	Not reported	No multivariate analysis on VTE was conducted.	Retrospective, no blinding
2012, Thibodeau et al. [67]	Cohort, VTE prophylaxis strategies uncertain	Cardiac transplantation (n=201)	Mean 1.8 to 3.6 years	Symptomatic DVT (12% in sirolimus treated vs 7% in nonsirolimus treated), PE (sirolimus 3% vs. nonsirolimus 0.7%), unclear modality	Not reported	VTE: Sirolimus treatment remained associated with an increased risk of VTE after adjusting for body mass index.	Retrospective, no blinding

Continued

Table 2. Continued

Study (Year of Publication, First Author, [Reference Number])	Type of Study and Use of LMWH, UFH, or Other Forms of VTE Prophylaxis	Type of Surgery and Sample Size	Duration of Follow-up	Incidence of Symptomatic DVT and PE, Diagnostic Modality	Incidence of Asymptomatic DVT and PE (%), Diagnostic Modality	Significant Adjusted Risk Factors for VTE and Complications	Bias Assessment
2014, Urbanski et al. [71]	Cohort, VTE prophylaxis strategies uncertain	Bentall procedure with either tissue or mechanical valve (n=29)	4.5 years	Symptomatic fatal PE (3.5%)	Not reported	No multivariate analysis on VTE was conducted.	Retrospective, no blinding
2011, Manliot et al. [72]	Cohort, heparinization of arterial and central venous catheters, antithrombotic therapy for aortopulmonary shunt or mechanical valve	Congenital heart disease requiring surgery (n=1542)	Until hospital discharge	Symptomatic DVT (7.9%), PE (0.3%), fatal PE (0.06%) by either US, CTPA, MRI, echocardiography, surgery, or postmortem	Not reported	VTE: 1. Younger than 31 days (OR, 2.0). 2. Oxygen saturation <85% (OR, 2.0). 3. Previous thrombosis (OR, 2.6). 4. Heart transplantation (OR, 4.1). 5. Deep hypothermic circulatory arrest (OR, 1.9). 6. Central venous catheter >5 days (OR 1.2). 7. Postoperative use of ventricular assist device or extracorporeal membrane oxygenation (OR, 5.2)	Retrospective, no blinding
2012, Manliot et al. [73]	Cohort, either without VTE, aspirin, or warfarin prophylaxis	SCPC (n=139) or Fontan (n=162)	Until Oct 2009 (since 2000–2009 after Fontan and since 2003–2008 for single ventricle cardiac lesions)	Symptomatic DVT after Fontan (21%) and after SCPC (34%) by radiology, surgery, catheterization findings, or postmortem, thrombotic complication as a cause of death (3.6%)	Not reported	VTE: 1. Use of thromboprophylaxis (HR, 0.2 after SCPC and HR, 0.27 warfarin vs. aspirin or 0.18 warfarin vs. none after Fontan)	Retrospective, no blinding
1995, Goldhaber et al. [66]	RCT, IPC with graduated compression stockings vs. graduated compression stockings alone. All patients received aspirin	CABG (n=344)	Until hospital discharge	Symptomatic PE (0.9%) by high probability V/Q scan, fatal PE (0.3%)	Asymptomatic DVT (21%) by routine USS	No multivariate analysis on VTE was conducted.	Prospective, no blinding

Continued

Table 2. Continued

Study (Year of Publication, First Author, [Reference Number])	Type of Study and Use of LMWH, UFH, or Other Forms of VTE Prophylaxis	Type of Surgery and Sample Size	Duration of Follow-up	Incidence of Symptomatic DVT and PE, Diagnostic Modality	Incidence of Asymptomatic DVT and PE (%), Diagnostic Modality	Significant Adjusted Risk Factors for VTE and Complications	Bias Assessment
1979, Pantely et al. [68]	RCT, Warfarin vs. control vs. high-dose antiplatelet agent group (aspirin 325 mg+dipyridamole 75 mg both 3 times per day)	CABG (n=65)	Until hospital discharge	Symptomatic PE (3%), DVT (1.5%), unclear modality	Not reported	No multivariate analysis on VTE was conducted.	Prospective, no blinding
2010, Schroeder et al. [69]	RCT, IV heparin (10 U/kg per hour vs. 5% dextrose) for 14 days or until central venous catheter removed	Congenital heart diseases (n=90)	Until day 14 or removal of central venous catheter	Not reported	Asymptomatic central venous catheter related thrombosis (16%) by echocardiography or USS	VTE: 1. Central venous catheter related thrombosis (16%) 2. Central venous catheter malfunction (OR, 11.2) Bleeding requiring interruption of study drug in 1 patient in the placebo group	Prospective, double blinded
2013, Weltert et al. [70]	RCT, VTE prophylaxis strategies uncertain	Mixed (n=1049)	Until hospital discharge	Symptomatic DVT (2.4%), unclear modality	Not reported	No multivariate analysis on VTE was conducted.	Prospective, no blinding
2013, Rahman et al. [75]	Cohort, enoxaparin 20 or 40 mg/day	Aortic surgery (n=189)	12 months after surgery	Not reported	Not reported	Incidence of pericardial effusion (21% after 40 mg enoxaparin vs. 19% after 20 mg enoxaparin). No multivariate analysis on pericardial effusion was conducted.	Retrospective, no blinding
1988, Ikaheimo et al. [76]	Cohort, warfarin from postoperative day 2 (valve surgery group 1), aspirin+dipyridamole on day of surgery (CABG group 2), or warfarin started 2 weeks before surgery (CABG group 3)	Mixed (n=150)	2 weeks after surgery	Not reported	Not reported	Incidence of pericardial effusion (68% group 1 vs. 80% group 2 vs. 84% group 3). No multivariate analysis on pericardial effusion was conducted.	Prospective, no blinding
2013, Alassar et al. [77]	Cohort, VTE prophylaxis strategies uncertain	Transcatheter aortic valve implantation (n=119)	4 years after surgery	Fatal PE (0.8%)	Not reported	No multivariate analysis on VTE outcome was conducted.	Retrospective, no blinding

Continued

Table 2. Continued

Study (Year of Publication, First Author, [Reference Number])	Type of Study and Use of LMWH, UFH, or Other Forms of VTE Prophylaxis	Type of Surgery and Sample Size	Duration of Follow-up	Incidence of Symptomatic DVT and PE, Diagnostic Modality	Incidence of Asymptomatic DVT and PE, Diagnostic Modality	Significant Adjusted Risk Factors for VTE and Complications	Bias Assessment
1982, McEnany et al. [78]	RCT, no UFH. Placebo vs aspirin 300 mg bd vs warfarin (INR, 1.5 to 2.0) starting on day 3 or 4 after surgery	CABG (n=216)	12 months after surgery	Symptomatic PE (1.9%)	Not reported	No multivariate analysis on VTE outcome was conducted, bleeding only occurred in 3 patients (4.4%) on warfarin.	Prospective, partial blinding for warfarin, double-blinded for aspirin group
2010, Garvin et al. [79]	Cohort, VTE prophylaxis strategies uncertain	CABG (n=1403)	Hospital discharge	Symptomatic PE (0.57%) by high probability V/Q scan or CTPA	Not reported	No multivariate analysis on VTE outcome was conducted.	Prospective, no blinding
2007, Jensen and Yang [80]	Cohort, VTE prophylaxis strategies uncertain	CABG (n=315)	Hospital discharge	Symptomatic PE (0.32%), unclear modality	Not reported	No multivariate analysis on VTE outcome was conducted.	Retrospective, no blinding
1975, Wisoff et al. [81]	Cohort, UFH for all patients	CABG (n=200)	Hospital discharge	Symptomatic PE (3.5%), unclear modality	Not reported	No multivariate analysis on VTE outcome was conducted.	Retrospective, no blinding
1994, Parenti [82]	Cohort, VTE prophylaxis strategies uncertain	CABG (n=120)	Hospital discharge	Symptomatic PE (4.1%)	Not reported	No multivariate analysis on VTE outcome was conducted.	Retrospective, no blinding
1991, Saito et al. [83]	Cohort, VTE prophylaxis strategies uncertain	CABG (n=8100)	Hospital discharge	Symptomatic PE (0.074%)	Not reported	No multivariate analysis on VTE outcome was conducted.	Retrospective, no blinding
1987, Dorros et al. [84]	Cohort, VTE prophylaxis strategies uncertain	CABG (n=674) >70 years old	1 year follow-up	Symptomatic PE (2.1%), unclear modality	Not reported	No multivariate analysis on VTE, bleeding requiring reoperation in 24 patients (3.6%), but uncertain whether bleeding was related to VTE prophylaxis as the latter information was not described.	Retrospective, no blinding

bd indicates twice a day; CABG, coronary artery bypass grafting; CHF, congestive heart failure; CTPA, computed tomographic pulmonary angiogram; DVT, deep vein thrombosis; GCS, graduate compression stockings; HIT, heparin-induced thrombocytopenia; HR, hazard ratio; ICD, International Classification of Diseases; IV, intravenous; INR, international normalized ratio; IPPV, invasive mechanical ventilation; LMWH, low-molecular-weight heparin; LVEF, left ventricular ejection fraction; MRI, magnetic resonance imaging; OR, odds ratio; PE, pulmonary embolism; RCT, randomized, controlled trial; SC, subcutaneous; CPC, superior cavaopulmonary connection; tds, three times a day; TEE, transesophageal echocardiography; UFH, unfractionated heparin; US, ultrasound scan; VTE, venous thromboembolism.




Figure 2. Forest plots showing the difference in risks of symptomatic or asymptomatic PE with and without using some forms of venous thromboembolism prophylaxis. ASP indicates aspirin; CI, confidence interval; IPC, intermittent pneumatic compression to the lower limbs; PE, pulmonary embolism; RR, relative risk; WARF, warfarin.

(RR, 3.0; 95% CI, 0.14–66; P=0.49). There were no RCTs directly comparing UFH or LMWH against aspirin alone for VTE prophylaxis after cardiac surgery.

Incidences and Risk Factors for Symptomatic and Asymptomatic VTE

The median incidence (interquartile range; IQR) of asymptomatic DVT and PE reported in the included studies were 14.8% (10.8–40.4) and 1.9% (0.4–25), respectively. The median incidence (IQR) of symptomatic DVT, PE, and fatal

PE were 3.2% (0.6–8.1), 0.6% (0.3–2.9), and 0.3% (0.08–1.7), respectively. After restricting our analysis to only prospective cohort studies or RCTs, the median incidence of symptomatic PE (0.7%; IQR, 0.1–1.7) and fatal PE (0.4%; IQR, 0.3–1.4) were similar to when all studies were considered. The incidence of symptomatic PE and fatal PE were not associated with the sample size of the included studies (P=0.606 and 0.142, respectively) or how recent the study was conducted (P=0.207 and 0.802, respectively). Incidence of VTE and modality used to diagnose DVT and PE for each RCT as well as observational study are described in Table 2. For prospective




Figure 3. Forest plots showing the difference in the risks of symptomatic and asymptomatic VTE with and without using VTE prophylaxis. ASP indicates aspirin; CI, confidence interval; HEP, unfractionated heparin; IPC, intermittent pneumatic compression to the lower limbs; LEG, leg elevation; RR, relative risk; VTE, venous thromboembolism; WARF, warfarin.




Figure 4. Benefits of VTE prophylaxis on risks of PE after excluding studies without adequate allocation concealment. ASP indicates aspirin; CI, confidence interval; IPC intermittent pneumatic compression to the lower limbs; PE, pulmonary embolism; RR, relative risk; VTE, venous thromboembolism; WARF, warfarin.

studies, all patients with clinical signs of DVT were expected to have an ultrasound scan (USS) as part of the studies, but the percentage of patients who had USS when clinical signs of DVT were present in the retrospective studies was not available.

We identified 2 large postmortem studies, and, in these 2 studies, fatal PE accounted for 11% to 20% of the unexplained deaths after cardiac surgery.^{57,58} Diagnosis of PE was suspected or diagnosed clinically by a CT pulmonary angiogram before death only in 33% to 42% of the patients.^{57,58} Many patients who died from fatal PE were misdiagnosed clinically as having acute heart failure, acute myocardial infarction, pneumonia, or sudden cardiac death.⁵⁷ Fatal PE was reported within 24 hours of cardiac surgery in at least 1 observational study.²⁶

The risk factors for VTE after cardiac surgery that remained significant after adjustment by a multivariate analysis are summarized in Table 3. Previous history of VTE, increasing age, obesity, left or right ventricular failure, prolonged bed rest/mechanical ventilation/use of a central venous catheter, and omission of all forms of anticoagulation or -platelet agents were the most consistent risk factors for VTE after cardiac surgery. Results of 2 meta-analyses on use of activated factor VIIa or tranexamic acid could not confirm these drugs as a significant risk factor for VTE after cardiac surgery.^{62,74} There was a suggestion that the leg with the great saphenous vein harvested could have a higher risk of DVT compared to the contralateral leg (RR, 1.48; 95% CI, 0.96–2.26; P=0.07; I²=71%), but this result did not reach statistical significance and had substantial heterogeneity




Figure 5. Benefits of VTE prophylaxis on risks of symptomatic VTE by excluding studies without adequate allocation concealment. ASP indicates aspirin; CI, confidence interval; HEP, unfractionated heparin; IPC, intermittent pneumatic compression to the lower limbs; LEG, leg elevation; RR, relative risk; VTE, venous thromboembolism; WARF, warfarin.




Figure 6. Funnel plot showing a lack of publication bias, using all forms of venous thromboembolic events as an outcome. RR indicates relative risk.

(Figure 7). Restricting the analysis to studies that used routine USS for both legs^{20,25,31,36} did not change the results significantly (RR, 1.58; 95% CI, 0.96–2.61; $P=0.07$; $I^2=77\%$). Metaregression suggested that sample size was likely the main reason behind this heterogeneity (slope=0.004; 95% CI, 0.002–0.006; $P=0.001$; Figure 8), with large sample size studies more likely to show an increased risk of DVT in the leg with the great saphenous vein harvested than smaller studies.

Incidences and Risk Factors for Complications From VTE Prophylaxis

Eight studies reported incidence of bleeding after cardiac surgery,^{17,19,25,53,54,78,84} but only 1 study reported risk factors for bleeding after multivariate analysis, with concurrent use of aspirin with systemic anticoagulation as the only significant risk factor in causing bleeding after valvular surgery, with some of these bleeding patients with an international normalized ratio (INR) >5 .⁵³

Although pericardial effusion was common after cardiac surgery (up to 84%),⁷⁶ using different doses of enoxaparin (20 vs. 40 mg/day) for VTE prophylaxis after proximal aortic surgery did not affect the incidence of pericardial effusion (19% vs. 21%, respectively) nor the risk of cardiac tamponade requiring surgical interventions.⁷⁵ Similarly, initiation of systemic anticoagulation for confirmed VTE also did not increase the risk of pericardial effusion in patients who took aspirin after CABG surgery.²⁷ Similar to bleeding after cardiac surgery, therapeutic systemic anticoagulation (INR >2.5) was the only reported risk factor for cardiac tamponade after valvular surgery in another observational study not included in this review.⁸⁵ No multivariate analysis on risk factors for cardiac tamponade or pericardial effusion was, however, presented in these studies.^{27,75,76,85}

Discussion

Cardiac surgery is increasingly offered to older patients with coexisting comorbidities,⁸⁶ and it is certain that the incidence of VTE after cardiac surgery will be increasing. Although VTE is an important preventable cause of morbidity and mortality in hospitalized patients, both the quality and amount of evidence to guide VTE prophylaxis after cardiac surgery were relatively sparse. With the evidence available, we confirmed that VTE prophylaxis could reduce the risk of PE and symptomatic VTE after cardiac surgery without increasing risk of bleeding and cardiac tamponade, unless the patients were systemically anticoagulated. Although some risk factors for VTE were specifically related to cardiac surgery, many important risk factors for VTE after cardiac surgery were indeed similar to risk factors for VTE in other patient populations. These results are clinically relevant and require further discussion.

First, although we demonstrated that some form of VTE prophylaxis was effective in reducing PE and symptomatic VTE, we could not identify whether 1 form of VTE prophylaxis was more effective than another, nor whether 1 particular pharmacological agent was superior to the others. Recent studies showed that platelets play a significant role in the pathogenesis of VTE and antiplatelet agents can have a protective effect against the first episode of VTE in patients with hip fracture or undergoing hip arthroplasty ($n=13\ 356$) and recurrent VTE after treatment with systemic anticoagulation.^{87–91} Our results showed that aspirin appears to be effective in reducing VTE after cardiac surgery in the pediatric population (using 5 mg/kg/day)⁴⁵ or when a higher dose of aspirin is used in adults (>300 mg/day).⁶⁸ Evidence suggests that aspirin resistance is common within the first week after cardiac surgery and the standard low-dose aspirin (100 mg/day) would not be adequate to exert its full antiplatelet effects in many patients.^{11–13} Because cardiac surgical patients are also prothrombotic postsurgery (up to 30 days),^{6–9} use of low-dose antiplatelet agents alone would not be sufficient in preventing VTE during this high-risk period if the patients have multiple risk factors for VTE (Table 3). We noted that fatal PE after cardiac surgery still occurred within the last decade despite advances in cardiac surgical care, including routine use of low-dose aspirin post-CABG.^{23,26,71,73,77} Fatal PE accounted for approximately 11% to 20% of all unexplained deaths after cardiac surgery, with at least 50% not diagnosed before death.^{57,58} Median incidence of fatal PE (0.3%) reported in this review was consistent with the data from these 2 postmortem reports if the overall mortality after cardiac surgery was approximately 3%, and this may, in part, explain why omission of early VTE prophylaxis is associated with an increased mortality in the critically ill.⁹² Initiating UFH or LMWH, which offers additional VTE protection above intermittent pneumatic lower-limb compression,^{93,94} should

Table 3. Risk Factors for VTE and Bleeding After Cardiac Surgery That Remained Significant After Adjustment by Multivariate Analysis in the Included Studies

	Reported Odds Ratio
Preoperative risk or protective factors for VTE	
Prior history of VTE	2.6, 3.1
Obesity	2.6 (or 1.2 per body mass index increment)
Charlson comorbidity index	1.2 (per index increment)
Increasing age	1.2 per 10 years increment
Female	NR
Previous right heart catheterization (or right heart failure)	2.9
Preoperative bed rest, acute surgery within 15 days of coronary angiography	2.8, 5.0, 6.0
Preoperative left ventricular failure (eg, ejection fraction <40%)	6.8
Pulmonary atresia with intact ventricular septum before Fontan surgery	3.6
Low bilirubin (suggestive of low right-sided filling pressure) before Fontan surgery	0.8
Cyanotic heart disease (oxygen saturation <85%)	2.0
Intraoperative risk factors for VTE	
Deep hypothermic circulatory arrest	1.9
Heart transplantation	4.1
Postoperative risk factors for VTE	
Left ventricular failure	4.1, 8.0
Bed rest >3 days	5.0
Mechanical ventilation >3 days	2.5 (or 1.02 per 10 hours ventilation)
Failed extubation requiring reintubation	2.6
Central venous catheter >7 to 10 days	4.3, 17.8
Presence of a malfunctioning central venous catheter	11.2
Transfusion	2.2, 2.8
Use of ventricular assist device or ECMO after congenital heart surgery	5.2
No heparin prophylaxis before discharge from surgical unit	NR
Subtherapeutic warfarin	3.5
No aspirin or warfarin after Fontan surgery	5.6
Use of sirolimus after cardiac transplantation	NR
Risk factors for bleeding	
Concomitant use of aspirin and systemic anticoagulation	7.4

ECMO indicates extracorporeal membrane oxygenation; NR, not reported; VTE, venous thromboembolism.

be seriously considered for all cardiac surgical patients as soon as possible, or on postoperative day 1, if the patients have no active bleeding, especially for those with multiple risk factors for VTE.

Second, we could not identify sufficient data to support the notion that use of low-dose UFH or LMWH would increase the risks of bleeding (eg, mediastinal, intracranial, or gastrointestinal hemorrhage), pericardial effusion, and cardiac tamponade.⁴ Though these complications are not rare after surgery,^{76,95,96} whether low-dose UFH or LMWH would

substantially increase such risks remains scientifically unproven. In one large, retrospective cohort study, use of UFH or LMWH at 48 hours post-CABG was not associated with an increased risk of bleeding compared to no VTE prophylaxis.¹⁹ Using different doses of enoxaparin (20 vs. 40 mg/day) for VTE prophylaxis after proximal aortic surgery also did not appear to affect incidence of pericardial effusion nor risk of cardiac tamponade requiring surgical interventions.⁷⁵ In fact, most studies reported that bleeding after cardiac surgery only occurred in the presence of systemic




Figure 7. Forest plot showing the incidence of deep vein thrombosis in the leg with great saphenous vein harvested against the leg without the great saphenous vein harvested for coronary artery bypass graft surgery. CI indicates confidence interval; LEG, leg elevation; RR, relative risk.

anticoagulation,^{35,52–54,78} especially when patients were overanticoagulated.^{53,54} Of all the studies included in this review, only 1 study reported risk factors for bleeding after using multivariate analysis, and it showed that only concurrent use of systemic anticoagulation with an antiplatelet agent was associated with an increased risk of bleeding.⁵³ Nevertheless, most reported studies are underpowered to detect small increases in rare side effects; clinicians should still be mindful about the rare, but serious, effect of pericardial hematoma before initiating anticoagulants. Perhaps, a lower than usual threshold to cease the bridging UFH (subcutaneously or intravenously) or LMWH would reduce a patient's risk of bleeding and cardiac tamponade, while the patient is taking concurrent aspirin and warfarin (eg, when INR >1.5).⁸⁵

The last consideration is the limitations of this systematic review. Although we have searched extensively for the evidence on risks and benefits of VTE prophylaxis after cardiac surgery, there is an obvious evidence gap, suggesting that an adequately powered RCT is needed to confirm whether initiating low-dose UFH or LMWH from postoperative day 1 after cardiac surgery would be more cost-effective than

using low- or high-dose aspirin, warfarin, or intermittent lower-limb pneumatic compression alone in reducing morbidity and mortality of VTE. Our results suggested that DVT could be more common in the leg with the great saphenous vein harvested for CABG. This finding is not surprising because the lower-limb venous system has been disrupted after the great saphenous vein is harvested. Whether this increased risk was also contributed by wound pain causing immobilization, intolerance to graduate compression stockings, or intermittent pneumatic compression remains uncertain. Finally, we noted that many studies have a relatively short follow-up period in detecting VTE. With those studies with an extended follow-up period beyond the acute hospital stay,^{23,44–46,49,59,71} symptomatic VTE continued to occur. Hence, future cohort studies and RCTs on VTE after cardiac surgery should consider VTE events well beyond the immediate postoperative period.

In conclusion, PE and symptomatic VTE are not rare after cardiac surgery and these events can be reduced with VTE prophylaxis. Data on whether one form of VTE prophylaxis is superior to another are sparse. Although VTE rates appeared to be similar between patients taking high-dose aspirin (>300 mg/day for adults and 5 mg/kg/day for children) and warfarin, low-dose aspirin (100 mg/day) used to keep the bypass graft patent is unlikely to be adequate in preventing VTE post-CABG, especially for patients with multiple risk factors for VTE. We found no evidence to support the notion that use of low-dose UFH or LMWH for VTE prophylaxis would increase risk of cardiac tamponade or bleeding after cardiac surgery. Bleeding after cardiac surgery is mainly related to systemic overanticoagulation or concurrent use of systemic anticoagulation and -platelet agents. Unless proven otherwise by adequately powered RCTs, initiating low-dose UFH or LMWH as soon as possible or on postoperative day 1 after cardiac surgery for patients who have no active bleeding is highly recommended, especially if they have multiple risk factors for VTE.




Figure 8. Meta-regression showing an association between sample size and differences in relative risk of having deep vein thrombosis in the leg with and without great saphenous veins harvested for coronary artery bypass surgery.

Acknowledgments

The authors thank Dr Mostafavi for providing unpublished data from his study on venous thromboembolism after cardiac surgery.

Sources of Funding

Dr Ho is funded by the Raine Medical Research Foundation and Western Australia Department of Health through the Raine Clinical Research Fellowship.

Disclosures

None of the authors have any conflict of interest to declare in relation to the materials presented in this study. The funding source has no influence on the content and the decision to publish this manuscript.

References

- Centers for Disease Control and Prevention (CDC). Venous thromboembolism in adult hospitalizations—United States, 2007–2009. *MMWR Morb Mortal Wkly Rep.* 2012;61:401–404.
- The burden of venous thromboembolism in Australia. Report by Access Economics Pty Limited for the Australian and New Zealand Working Party on the management and prevention of venous thromboembolism in Australia. 2008 May. Available at: https://www.google.com.au/?gws_rd=ssl#q=The+burden+of+venous+thromboembolism+in+Australia.+Report+by+Access. Accessed October 15, 2015.
- Dunning J, Versteegh M, Fabbri A, Pavie A, Kolb P, Lockowandt U, Nashef SA; EACTS Audit and Guidelines Committee. Guideline on antiplatelet and anticoagulation management in cardiac surgery. *Eur J Cardiothorac Surg.* 2008;34:73–92.
- Cikirkcioglu M, Myers PO, Kalangos A. First do no harm: postoperative thromboprophylaxis following open heart surgery. *Eur J Cardiothorac Surg.* 2013;44:e184.
- Gould MK, Garcia DA, Wren SM, Karanicolas PJ, Arcelus JI, Heit JA, Samama CM; American College of Chest Physicians. Prevention of VTE in nonorthopedic surgical patients: antithrombotic therapy and prevention of thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest.* 2012;141:e227S–e277S.
- Parolari A, Mussoni L, Frigerio M, Naliato M, Alamanni F, Galanti A, Fiore G, Veglia F, Tremoli E, Bigliali P, Camera M. Increased prothrombotic state lasting as long as one month after on-pump and off-pump coronary surgery. *J Thorac Cardiovasc Surg.* 2005;130:303–308.
- Lison S, Dietrich W, Braun S, Boehm J, Schuster T, Englhard A, Perchuc A, Spannagl M, Busley R. Enhanced thrombin generation after cardiopulmonary bypass surgery. *Anesth Analg.* 2011;112:37–45.
- Vallely MP, Bannon PG, Bayfield MS, Hughes CF, Kriparides L. Quantitative and temporal differences in coagulation, fibrinolysis and platelet activation after on-pump and off-pump coronary artery bypass surgery. *Heart Lung Circ.* 2009;18:123–130.
- Edelman JJ, Reddel CJ, Kriparides L, Bannon PG, Fraser JF, Curnow JL, Vallely MP. Natural history of hypercoagulability in patients undergoing coronary revascularization and effect of preoperative myocardial infarction. *J Thorac Cardiovasc Surg.* 2014;148:536–543.
- Romlin BS, Söderlund F, Wählander H, Nilsson B, Baghaei F, Jeppsson A. Platelet count and function in paediatric cardiac surgery: a prospective observational study. *Br J Anaesth.* 2014;113:847–854.
- Brambilla M, Parolari A, Camera M, Colli S, Eligini S, Centenaro C, Anselmo A, Alamanni F, Tremoli E. Effect of two doses of aspirin on thromboxane biosynthesis and platelet function in patients undergoing coronary surgery. *Thromb Haemost.* 2010;103:516–524.
- Zimmermann N, Wenk A, Kim U, Kienzle P, Weber AA, Gams E, Schröer K, Hohlfeld T. Functional and biochemical evaluation of platelet aspirin resistance after coronary artery bypass surgery. *Circulation.* 2003;108:542–547.
- Emani S, Trainor B, Zurkowski D, Baird CW, Flynn-Thompson FE, Pigula FA, Emani SM. Aspirin unresponsiveness predicts thrombosis in high-risk pediatric patients after cardiac surgery. *J Thorac Cardiovasc Surg.* 2014;148:810–814.
- Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ.* 2003;327:557–560.
- Ramos RS, Salem BI, Haikal M, Gowda S, Coordes C, Leidenfrost R. Critical role of pulmonary angiography in the diagnosis of pulmonary emboli following cardiac surgery. *Cathet Cardiovasc Diagn.* 1995;36:112–117.
- Ramos R, Salem BI, De Pawlikowski MP, Coordes C, Eisenberg S, Leidenfrost R. The efficacy of pneumatic compression stockings in the prevention of pulmonary embolism after cardiac surgery. *Chest.* 1996;109:82–85.
- Mirhosseini SJ, Forouzannia SK, Mostafavi Pour Manshadi SM, Ali-Hassan-Sayegh S, Naderi N, Sanatkar M. Comparison of aspirin plus heparin with heparin alone on asymptomatic perioperative deep vein thrombosis in candidates for elective off-pump coronary artery bypass graft: a randomized clinical trial. *Cardiol J.* 2013;20:139–143.
- Pouplard C, May MA, Lochmann S, Amiral J, Vissac AM, Marchand M, Gruel Y. Antibodies to platelet factor 4-heparin after cardiopulmonary bypass in patients anticoagulated with unfractionated heparin or a low-molecular-weight heparin: clinical implications for heparin-induced thrombocytopenia. *Circulation.* 1999;99:2530–2536.
- Kulik A, Rassen JA, Myers J, Schneeweiss S, Gagne J, Polinski JM, Liu J, Fischer MA, Choudry NK. Comparative effectiveness of preventative therapy for venous thromboembolism after coronary artery bypass graft surgery. *Circ Cardiovasc Interv.* 2012;5:590–596.
- Ambrosetti M, Salerno M, Zambelli M, Mastropasqua F, Tramari R, Pedretti RF. Deep vein thrombosis among patients entering cardiac rehabilitation after coronary artery bypass surgery. *Chest.* 2004;125:191–196.
- Frizzelli R, Tortelli O, Di Comite V, Ghirardi R, Pinzi C, Scarduelli C. Deep venous thrombosis of the neck and pulmonary embolism in patients with a central venous catheter admitted to cardiac rehabilitation after cardiac surgery: a prospective study of 815 patients. *Intern Emerg Med.* 2008;3:325–330.
- DeLaria GA, Hunter JA. Deep venous thrombosis. Implications after open heart surgery. *Chest.* 1991;99:284–288.
- Ho KM, Bertenshaw C, Same S, Schneider M, Williams KA, Godsell T, Hird K. Differential associations between body mass index and outcomes after elective adult cardiac surgery: a linked data cohort study. *Anaesth Intensive Care.* 2013;41:573–583.
- Hanson EC, Levine FH. Hyperlipoproteinemia as a significant risk factor for pulmonary embolism in patients undergoing coronary artery bypass grafting. *Ann Thorac Surg.* 1982;33:593–598.
- Schwann TA, Kistler L, Engoren MC, Habib RH. Incidence and predictors of postoperative deep vein thrombosis in cardiac surgery in the era of aggressive thromboprophylaxis. *Ann Thorac Surg.* 2010;90:760–766.
- Saranteas T, Kostopanagiotou G, Tzoufi M, Drachidi K, Knox GM, Panou F. Incidence of inferior vena cava thrombosis detected by transthoracic echocardiography in the immediate postoperative period after adult cardiac and general surgery. *Anaesth Intensive Care.* 2013;41:782–787.
- Ambrosetti M, Ageno W, Salerno M, Pedretti RF. Postoperative pericardial effusion in patients receiving anticoagulants for deep vein thrombosis after coronary artery bypass graft surgery. *J Thromb Haemost.* 2005;3:2367–2368.
- Rao G, Zikria EA, Miller WH, Samadani SR, Ford WB. Incidence and prevention of pulmonary embolism after coronary artery surgery. *Vasc Surg.* 1975;9:37–45.
- Josa M, Siouffi SY, Silverman AB, Barsamian EM, Khuri SF, Sharma GV. Pulmonary embolism after cardiac surgery. *J Am Coll Cardiol.* 1993;21:990–996.
- Gillinov AM, Davis EA, Alberg AJ, Rykiel M, Gardner TJ, Cameron DE. Pulmonary embolism in the cardiac surgical patient. *Ann Thorac Surg.* 1992;53:988–991.
- Reis SE, Polak JF, Hirsch DR, Cohn LH, Creager MA, Donovan BC, Goldhaber SZ. Frequency of deep venous thrombosis in asymptomatic patients with coronary artery bypass grafts. *Am Heart J.* 1991;122:478–482.
- Egawa N, Hiromatsu S, Shintani Y, Kanaya K, Fukunaga S, Aoyagi S. Prevention of venous thromboembolism in thoracic and cardiovascular surgery. *Asian Cardiovasc Thorac Ann.* 2009;17:505–509.
- Lahtinen J, Ahvenjärvi L, Biancari F, Ojala R, Mosorin M, Cresti R, Lepojärvi M, Juvinen T. Pulmonary embolism after off-pump coronary artery bypass surgery as detected by computed tomography. *Am J Surg.* 2006;192:396–398.
- Mitra B, Phillips L, Cameron PA, Billah B, Reid C. The safety of recombinant factor VIIa in cardiac surgery. *Anaesth Intensive Care.* 2010;38:671–677.
- Canver CC, Fiedler RC. Venous thromboembolic complications after open heart surgery. *Vasc Endovascular Surg.* 1992;26:213–217.

36. Ayatollahzade-Isfahani F, Pashang M, Omran AS, Saadat S, Shirani S, Fathollahi MS. Comparing the impact of supine and leg elevation positions during coronary artery bypass graft on deep vein thrombosis occurrence: a randomized clinical trial study. *J Vasc Nurs.* 2013;31:64–67.
37. Gandhi MJ, Pierce RA, Zhang L, Moon MR, Despotis GJ, Moazami N. Use of activated recombinant factor VII for severe coagulopathy post ventricular assist device or orthotopic heart transplant. *J Cardiothorac Surg.* 2007;2:32.
38. Mueller J, Jeudy J, Poston R, White CS. Cardiac CT angiography after coronary bypass surgery: prevalence of incidental findings. *AJR Am J Roentgenol.* 2007;189:414–419.
39. Nussmeier NA, Whelton AA, Brown MT, Langford RM, Hoeft A, Parlow JL, Boyce SW, Verburg KM. Complications of the COX-2 inhibitors parecoxib and valdecoxib after cardiac surgery. *N Engl J Med.* 2005;352:1081–1091.
40. Belczak CE, Tyszka AL, Godoy JM, Ramos RN, Belczak SQ, Caffaro RA. Clinical complications of limb undergone harvesting of great saphenous vein for coronary artery bypass grafting using bridge technique. *Rev Bras Cir Cardiovasc.* 2009;24:68–72.
41. ElBardissi AW, Shekar P, Couper GS, Cohn LH. Minimally invasive aortic valve replacement in octogenarian, high-risk, transcatheter aortic valve implantation candidates. *J Thorac Cardiovasc Surg.* 2011;141:328–335.
42. Gill R, Herbertson M, Vuylsteke A, Olsen PS, von Heymann C, Mythen M, Sellke F, Booth F, Schmidt TA. Safety and efficacy of recombinant activated factor VII: a randomized placebo-controlled trial in the setting of bleeding after cardiac surgery. *Circulation.* 2009;120:21–27.
43. Alfirevic A, Duncan A, You J, Lober C, Soltesz E. Recombinant factor VII is associated with worse survival in complex cardiac surgical patients. *Ann Thorac Surg.* 2014;98:618–624.
44. White RH, Zhou H, Romano PS. Incidence of symptomatic venous thromboembolism after different elective or urgent surgical procedures. *Thromb Haemost.* 2003;90:446–455.
45. McCrindle BW, Manlhiot C, Cochrane A, Roberts R, Hughes M, Szekhtman B, Weintraub R, Andrew M, Monagle P; Fontan Anticoagulation Study Group. Factors associated with thrombotic complications after the Fontan procedure: a secondary analysis of a multicenter, randomized trial of primary thrombo-prophylaxis for 2 years after the Fontan procedure. *J Am Coll Cardiol.* 2013;61:346–353.
46. Zahn R, Gerckens U, Linke A, Sievert H, Kahlert P, Hambrecht R, Sack S, Abdel-Wahab M, Hoffmann E, Schiele R, Schneider S, Senges J; German Transcatheter Aortic Valve Interventions-Registry Investigators. Predictors of one-year mortality after transcatheter aortic valve implantation for severe symptomatic aortic stenosis. *Am J Cardiol.* 2013;112:272–279.
47. Iribarne A, Burgener JD, Hong K, Raman J, Akhter S, Easterwood R, Jeevanandam V, Russo MJ. Quantifying the incremental cost of complications associated with mitral valve surgery in the United States. *J Thorac Cardiovasc Surg.* 2012;143:864–872.
48. Thielmann M, Bunschkowski M, Tossios P, Selleng S, Marggraf G, Greinacher A, Jakob H, Massoudy P. Perioperative thrombocytopenia in cardiac surgical patients—incidence of heparin-induced thrombocytopenia, morbidities and mortality. *Eur J Cardiothorac Surg.* 2010;37:1391–1395.
49. Rosenthal DN, Friedman AH, Kleinman CS, Kopf GS, Rosenfeld LE, Hellenbrand WE. Thromboembolic complications after Fontan operations. *Circulation.* 1995;92:II287-II293.
50. Michalopoulos A, Tzelepis G, Dafni U, Geroulanos S. Determinants of hospital mortality after coronary artery bypass grafting. *Chest.* 1999;115:1598–1603.
51. Briffa N, Large SR. Venous thromboembolism in coronary artery surgery. *Heart.* 1996;75:433.
52. Doss M, Wood JP, Kiesling AH, Moritz A. Comparative evaluation of left ventricular mass regression after aortic valve replacement: a prospective randomized analysis. *J Cardiothorac Surg.* 2011;6:136.
53. Bucci C, Geerts WH, Sinclair A, Fremes SE. Comparison of the effectiveness and safety of low-molecular weight heparin versus unfractionated heparin anticoagulation after heart valve surgery. *Am J Cardiol.* 2011;107:591–594.
54. Montalescot G, Polle V, Collet JP, Leprince P, Bellanger A, Gandjbakhch I, Thomas D. Low molecular weight heparin after mechanical heart valve replacement. *Circulation.* 2000;101:1083–1086.
55. Lee JD, Lee SJ, Tsushima WT, Yamauchi H, Lau WT, Popper J, Stein A, Johnson D, Lee D, Petrovitch H, Dang CR. Benefits of off-pump bypass on neurologic and clinical morbidity: a prospective randomized trial. *Ann Thorac Surg.* 2003;76:18–25.
56. Riess FC, Poetzsch B, Madlener K, Cramer E, Doll KN, Doll S, Lorke DE, Kormann J, Mueller-Berghaus G. Recombinant hirudin for cardiopulmonary bypass anticoagulation: a randomized, prospective, and heparin-controlled pilot study. *Thorac Cardiovasc Surg.* 2007;55:233–238.
57. Rastan AJ, Gummert JF, Lachmann N, Walther T, Schmitt DV, Falk V, Doll N, Caffier P, Richter MM, Wittekind C, Mohr FW. Significant value of autopsy for quality management in cardiac surgery. *J Thorac Cardiovasc Surg.* 2005;129:1292–1300.
58. Zehr KJ, Liddicoat JR, Salazar JD, Gillinov AM, Hruban RH, Hutchins GM, Cameron DE. The autopsy: still important in cardiac surgery. *Ann Thorac Surg.* 1997;64:380–383.
59. Heard BE. Pathology of hearts after aortocoronary saphenous vein bypass grafting for coronary artery disease, studied by post-mortem coronary angiography. *Br Heart J.* 1976;38:838–859.
60. Song HK, Tibayan FA, Kahl EA, Sera VA, Slater MS, Deloughery TG, Scanlan MM. Safety and efficacy of prothrombin complex concentrates for the treatment of coagulopathy after cardiac surgery. *J Thorac Cardiovasc Surg.* 2014;147:1036–1040.
61. Sanchez R, Haft JL. Temporal relationship of complications after coronary artery bypass graft surgery: scheduling for safe discharge. *Am Heart J.* 1994;127:282–286.
62. Adler Ma SC, Brindle W, Burton G, Gallacher S, Hong FC, Manelius I, Smith A, Ho W, Alston RP, Bhattacharya K. Tranexamic acid is associated with less blood transfusion in off-pump coronary artery bypass graft surgery: a systematic review and meta-analysis. *J Cardiothorac Vasc Anesth.* 2011;25:26–35.
63. Marrone C, Galasso G, Piccolo R, de Leva F, Paladini R, Piscione F, Santoro G. Antiplatelet versus anticoagulation therapy after extracardiac conduit Fontan: a systematic review and meta-analysis. *Pediatr Cardiol.* 2011;32:32–39.
64. Beghi C, Fragnito C, Antonelli A, Reverberi C, Ferrari P, Saccani S, Fesani F. Prevention of deep venous thrombosis by a new low molecular weight heparin (Fluxum) in cardiac surgery. *Int Angiol.* 1993;12:383–386.
65. Clark KB, Kon ND, Hammon JW Jr, MacGregor DA, Kincaid EH, Reichert MG. Factor IX complex for the treatment of severe bleeding after cardiac surgery. *J Cardiovasc Pharmacol.* 2013;62:67–71.
66. Goldhaber SZ, Hirsch DR, MacDougall RC, Polak JF, Creager MA, Cohn LH. Prevention of venous thrombosis after coronary artery bypass surgery (a randomized trial comparing two mechanical prophylaxis strategies). *Am J Cardiol.* 1995;76:993–996.
67. Thibodeau JT, Mishkin JD, Patel PC, Kaiser PA, Ayers CR, Mammen PP, Markham DW, Ring WS, Peltz M, Drazner MH. Sirolimus use and incidence of venous thromboembolism in cardiac transplant recipients. *Clin Transplant.* 2012;26:953–959.
68. Pantely GA, Goodnight SH Jr, Rahimtoola SH, Harlan BJ, DeMots H, Calvin L, Rösch J. Failure of antiplatelet and anticoagulant therapy to improve patency of grafts after coronary-artery bypass: a controlled, randomized study. *N Engl J Med.* 1979;301:962–966.
69. Schroeder AR, Axelrod DM, Silverman NH, Rubesova E, Merkel E, Roth SJ. A continuous heparin infusion does not prevent catheter-related thrombosis in infants after cardiac surgery. *Pediatr Crit Care Med.* 2010;11:489–495.
70. Weltz L, Nardella S, Rondinelli MB, Pierelli L, De Paulis R. Reduction of allogeneic red blood cell usage during cardiac surgery by an integrated intra- and postoperative blood salvage strategy: results of a randomized comparison. *Transfusion.* 2013;53:790–797.
71. Urbanski PP, Dinstak W, Rents W, Heinz N, Diegeler A. Long-term results after aortic root replacement using self-assembled valve composite grafts in patients with small aortic annulus. *Interact Cardiovasc Thorac Surg.* 2014;18:159–163.
72. Manlhiot C, Menjak IB, Brandão LR, Gruenwald CE, Schwartz SM, Sivarajan VB, Yoon H, Maratta R, Carew CL, McMullen JA, Clarizia NA, Holtby HM, Williams S, Calderone CA, Van Arsdell GS, Chan AK, McCrindle BW. Risk, clinical features, and outcomes of thrombosis associated with pediatric cardiac surgery. *Circulation.* 2011;124:1511–1519.
73. Manlhiot C, Brandão LR, Kwok J, Kegel S, Menjak IB, Carew CL, Chan AK, Schwartz SM, Sivarajan VB, Calderone CA, Van Arsdell GS, McCrindle BW. Thrombotic complications and thromboprophylaxis across all three stages of single ventricle heart palliation. *J Pediatr.* 2012;161:513–519.e3.
74. Ponschab M, Landoni G, Biondi-Zoccai G, Bignami E, Frati E, Nicolotti D, Monaco F, Pappalardo F, Zangrillo A. Recombinant activated factor VII increases stroke in cardiac surgery: a meta-analysis. *J Cardiothorac Vasc Anesth.* 2011;25:804–810.
75. Rahman IA, Hussain A, Davies A, Bryan AJ. NICE thromboprophylaxis guidelines are not associated with increased pericardial effusion after surgery of the proximal thoracic aorta. *Ann R Coll Surg Engl.* 2013;95:433–436.
76. Ikäheimo MJ, Huikuri HV, Airaksinen KE, Korhonen UR, Linnaluoto MK, Tarkka MR, Takkunen JT. Pericardial effusion after cardiac surgery: incidence, relation to the type of surgery, antithrombotic therapy, and early coronary bypass graft patency. *Am Heart J.* 1988;116:97–102.

77. Alassar A, Roy D, Valencia O, Brecker S, Jahangiri M. Survival, predictive factors, and causes of mortality following transcatheter aortic valve implantation. *Innovations (Phila)*. 2013;8:359–363.
78. McEnany MT, Salzman EW, Mundth ED, DeSanctis RW, Harthorne JW, Weintraub RM, Gates S, Austen WG. The effect of antithrombotic therapy on patency rates of saphenous vein coronary artery bypass grafts. *J Thorac Cardiovasc Surg*. 1982;83:81–89.
79. Garvin S, Muehlschlegel JD, Perry TE, Chen J, Liu KY, Fox AA, Collard CD, Aranki SF, Shernan SK, Body SC. Postoperative activity, but not preoperative activity, of antithrombin is associated with major adverse cardiac events after coronary artery bypass graft surgery. *Anesth Analg*. 2010;111:862–869.
80. Jensen L, Yang L. Risk factors for postoperative pulmonary complications in coronary artery bypass graft surgery patients. *Eur J Cardiovasc Nurs*. 2007;6:241–246.
81. Wisoff BG, Hartstein ML, Aintablian A, Hamby RL. Risk of coronary surgery. Two hundred consecutive patients with no hospital deaths. *J Thorac Cardiovasc Surg*. 1975;69:669–673.
82. Parenti C. Pulmonary embolism after coronary artery bypass surgery. *Crit Care Nurs Q*. 1994;17:48–50.
83. Saito T, Terada Y, Fukuda S, Sakao Y, Takayama T, Suma H, Wanibuchi Y, Furuta S. Acute pulmonary thromboembolism after coronary artery bypass grafting. *Kyobu Geka*. 1991;44:1003–1005.
84. Dorros G, Lewin RF, Daley P, Assa J. Coronary artery bypass surgery in patients over age 70 years: report from the Milwaukee Cardiovascular Data Registry. *Clin Cardiol*. 1987;10:377–382.
85. Malouf JF, Alam S, Gharzeddine W, Stefadouros MA. The role of anticoagulation in the development of pericardial effusion and late tamponade after cardiac surgery. *Eur Heart J*. 1993;14:1451–1457.
86. Ho KM. Determinants of the relationship between cost and survival time after elective adult cardiac surgery. *Anaesth Intensive Care*. 2014;42:303–309.
87. Prevention of pulmonary embolism and deep vein thrombosis with low dose aspirin: pulmonary embolism prevention (PEP) trial. *Lancet*. 2000;355:1295–1302.
88. Becker RC. Aspirin and the prevention of venous thromboembolism. *N Engl J Med*. 2012;366:2028–2030.
89. Ho KM, Yip CB, Duff O. Reactive thrombocytosis and risk of subsequent venous thromboembolism: a cohort study. *J Thromb Haemost*. 2012;10:1768–1774.
90. Ho KM, Chavan S. Prevalence of thrombocytosis in critically ill patients and its association with symptomatic acute pulmonary embolism. A multicentre registry study. *Thromb Haemost*. 2013;109:272–279.
91. Becattini C, Agnelli G, Schenone A, Eichinger S, Bucherini E, Silingardi M, Bianchi M, Moia M, Ageno W, Vandelli MR, Grandone E, Prandoni P; WARFASA Investigators. Aspirin for preventing the recurrence of venous thromboembolism. *N Engl J Med*. 2012;366:1959–1967.
92. Ho KM, Chavan S, Pilcher D. Omission of early thromboprophylaxis and mortality in critically ill patients: a multicenter registry study. *Chest*. 2011;140:1436–1446.
93. Ho KM, Tan JA. Stratified meta-analysis of intermittent pneumatic compression of the lower limbs to prevent venous thromboembolism in hospitalized patients. *Circulation*. 2013;128:1003–1020.
94. Ho KM. Benefit of intermittent pneumatic compression of lower limbs in reducing venous thromboembolism in hospitalised patients: interactions between risk and effectiveness. *Anaesth Intensive Care*. 2014;42:140–141.
95. Pepi M, Muratori M, Barbier P, Doria E, Arena V, Berti M, Celeste F, Guazzi M, Tamborini G. Pericardial effusion after cardiac surgery: incidence, site, size, and haemodynamic consequences. *Br Heart J*. 1994;72:327–331.
96. Pompilio G, Filippini S, Agrifoglio M, Merati E, Lauri G, Salis S, Almanni F, Parolari A. Determinants of pericardial drainage for cardiac tamponade following cardiac surgery. *Eur J Cardiothorac Surg*. 2011;39:e107–e113.

Incidence of Venous Thromboembolism and Benefits and Risks of Thromboprophylaxis After Cardiac Surgery: A Systematic Review and Meta-Analysis

Kwok M. Ho, Ebrahim Bham and Warren Pavey

J Am Heart Assoc. 2015;4:e002652; originally published October 26, 2015;
doi: 10.1161/JAHA.115.002652

The *Journal of the American Heart Association* is published by the American Heart Association, 7272 Greenville Avenue,
Dallas, TX 75231
Online ISSN: 2047-9980

The online version of this article, along with updated information and services, is located on the World Wide Web at:

<http://jaha.ahajournals.org/content/4/10/e002652>

Subscriptions, Permissions, and Reprints: The *Journal of the American Heart Association* is an online only Open Access publication. Visit the Journal at <http://jaha.ahajournals.org> for more information.